

Ammar Anbari, MD

Shoulder, Hip and Knee Arthroscopy and Reconstruction www.CTSportsDoc.com Office Phone: (860) 889-7345

ROTATOR CUFF REPAIR PRESCRIPTION (includes subscapularis repair)

Date
Diagnosis
Date of Surgery
Frequency: 1 2 3 4 times/week Duration: 1 2 3 4 5 6 Weeks
Weeks 1-8: • True PROM only! The rotator cuff tendon needs to heal back into the bone *** LIMIT ER TO 30 degrees ONLY FOR 4 WEEKS FOR SUBSCAPULARIS • ROM goals: 140° FF/30° ER at side; ABD max 60° without rotation • No resisted motions of shoulder until 12 weeks post-op • Grip strengthening • No canes/pulleys until 6 weeks post-op, because these are active-assist exercises • Heat before PT, ice after PT
 Weeks 8-12: Begin AAROM → AROM as tolerated Goals: Same as above, but can increase as tolerated Light passive stretching at end ranges No strengthening/resisted motions yet, except grip strengthening Isometrics with arm at side beginning at 8 weeks
Months 3-12: • Advance to full ROM as tolerated with passive stretching at end ranges • Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers • Only do strengthening 3x/week to avoid rotator cuff tendonitis • Begin eccentrically resisted motions, plyometrics (ex. Weighted ball toss), proprioception (es. body blade) • Begin sports related rehab at 4 ½ months, including advanced conditioning • Return to throwing at 6 months • Throw from pitcher's mound at 9 months • Collision sports at 9 mohts • MMI is usually at 12 months post-op
X Modalities _X_ Electric StimulationX_ Ultrasound
Signature:Ammar Anbari, MD