

## Ammar Anbari, MD

Shoulder, Hip and Knee Arthroscopy and Reconstruction www.CTSportsDoc.com Office Phone: (860) 889-7345

## **ROTATOR CUFF REPAIR PRESCRIPTION**

Name
Date
Diagnosis
Date of Surgery
Frequency: 1 2 3 4 times/week Duration: 1 2 3 4 5 6 Weeks
Weeks 1-6:  • True PROM only! The rotator cuff tendon needs to heal back into the bone  • ROM goals: 140° FF/40° ER at side; ABD max 60-80° without rotation  • No resisted motions of shoulder until 12 weeks post-op  • Grip strengthening  • No canes/pulleys until 6 weeks post-op, because these are active-assist exercises  • Heat before PT, ice after PT
Weeks 6-12:  • Begin AAROM → AROM as tolerated  • Goals: Same as above, but can increase as tolerated  • Light passive stretching at end ranges  • No strengthening/resisted motions yet, except grip strengthening  • Isometrics with arm at side beginning at 8 weeks
Months 3-12:  • Advance to full ROM as tolerated with passive stretching at end ranges • Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilizers • Only do strengthening 3x/week to avoid rotator cuff tendonitis • Begin eccentrically resisted motions, plyometrics (ex. Weighted ball toss), proprioception (es. body blade) • Begin sports related rehab at 4 ½ months, including advanced conditioning • Return to throwing at 6 months • Throw from pitcher's mound at 9 months • Collision sports at 9 months • MMI is usually at 12 months post-op
_X_ Modalities _X_ Electric Stimulation
Signature:Ammar Anbari, MD