

## Ammar Anbari, MD

Shoulder, Hip and Knee Arthroscopy and Reconstruction www.CTSportsDoc.com Office Phone: (860) 889-7345

## **Subacromial Decompression Prescription**

Name
Date
Diagnosis
Date of Surgery
Frequency: 1 2 3 4 times/week Duration: 1 2 3 4 5 6 Weeks
Weeks 1-4:  • PROM → AAROM → AROM as tolerated  • ROM goals: 140° FF/40° ER at side  • No abduction-rotation until 4-8 weeks post-op  • No resisted motions until 4 weeks post-op  • D/C sling at 1-2 weeks post-op; sling only when sleeping if needed  • Heat before/ice after PT sessions
Weeks 4-8:  • D/C sling totally if not done previously  • Increase AROM in all directions with passive stretching at end ranges to maintain shoulder flexibility  • Goals: 160° FF/60° ER at side  • Begin light isometrics with arm at side for rotator cuff and deltoid; can advance to bands as tolerated  • Physical modalities per PT discretion
Weeks 8-12:  • Advance strengthening as tolerated: isometrics → bands → weights; 10 reps/1 set per rotator cuff, deltoid, and scapular stabilizers  • Only do strengthening 3x/week to avoid rotator cuff tendonitis  • If ROM lacking, increase to full with passive stretching at end ranges  • Begin eccentrically resisted motions, plyometrics, and closed chain exercises.
_X_ Modalities _X_ Electric Stimulation _X_ Ultrasound
Signature:Ammar Anbari, MD
/ \lillia / \libai, wb